

APPLICATION OF BIOFLOC TECHNOLOGY IN INTENSIVE FARMING AFFECTED PRODUCTION AND BLOOD PERFORMANCES OF THE CATF

by Sri Hastuti

Submission date: 27-Aug-2018 10:37 AM (UTC+0700)

Submission ID: 993574999

File name: 201602-BIOFLOC_TECHNOLOGY__Internasional.pdf (366.89K)

Word count: 4184

Character count: 21322

APPLICATION OF BIOFLOC TECHNOLOGY IN INTENSIVE FARMING AFFECTED PRODUCTION AND BLOOD PERFORMANCES OF THE CATFISH [*Clarias gariepinus* (Burchell, 1822)]

Sri Hastuti*, Subandiyono

Lab. of Aquaculture, Faculty of Fisheries and Marine Science,
Diponegoro University, Jl. Prof. Soedarto, SH Tembalang,
Semarang – 50275, Indonesia

Article history

Received

4 November 2015

Received in revised form


4 December 2015

Accepted

5 February 2016

*Corresponding author
hastuti_hastuti@yahoo.com

Graphical abstract



Abstract

The biofloc technology can control water quality under negligible water exchange. The aim of this study was to evaluate the effects of stocking density on production and blood performance of catfish (*Clarias gariepinus* [Burchell, 1822]). The catfish were reared in biofloc system (heterotrophic bacteria and addition of sugar for a period of 32 wk and used three levels of density i.e. (500, 1 000, and 1 500) fish per m². The production decreased with the increasing of stocking density, relative growth rate decreased with the increasing of that. The stocking density of 1 500 fish per m² on the biofloc technology can support maximum catfish production, health and proper water quality.

Keywords: Biofloc technology, catfish [*Clarias gariepinus* (Burchell, 1822)], growth, intensive, production

© 2016 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

The African catfish (*Clarias gariepinus* [Burchell, 1822]) could be cultured at high density up to 500 kg m⁻³, with specific growth rate value were 3.8 %, and food conversion rate (FCR) 0.8 g g⁻¹. Stocking density of fish might affect various aspects of the health of farmed fish. The healthy of catfish, *Clarias gariepinus*, was influenced by growth cycle. The welfare indicators were growth performance, behavior and physiological condition [1]. Production, growth and survival of that fish are known to be influenced by stocking density [2]. High stocking density in commercial catfish aquaculture system can increase fish production [1, 2].

Intensive catfish aquaculture system can lead to ammonification from feed residue and fish excretion, in which organic waste and ammonia can be easily accumulated in water which can reach an unsafe level. High concentration of ammonia is toxic to most teleost. Ammonia in waters

can be removed by biofloc system. The basic principle of biofloc technology is the retention of waste and its conversion to biofloc as a natural food within the culture system. This is done by constant aeration and agitation of water column and addition of carbon sources as organic substrate to allow aerobic decomposition and maintain high level of microbial floc in suspension in fed and or fertilized ponds [3-5]. The C: N ratio was balance of carbon to nitrogen as nutrient for develop and regeneration of the biofloc in the culture tanks. Theoretically, increased C:N ratio through carbon addition enhances conversion of toxic inorganic nitrogen to microbial biomass available as food for culture catfish. The optimum C:N ratio in an aquaculture can be maintained by adding carbon source or reduction of protein contain in feed [6].

The basic principles of activated suspension technique (AST), recently referred to as biofloc technology (BFT), is the retention of waste and its conversion to biofloc as a natural food within the

culture system. Biofloc technology has been widely applied in aquaculture. This system is not only an adequate approach in maintaining water quality in aquaculture system but also generate biomass that can contribute as a protein source for the cultured fish *in situ* [7]. Biofloc technology is useful for contributed to the growth and production of fish and it contains a good nutritional quality for herbivorous and omnivorous fish [5]. The principle of this system is assimilation of excreted dissolved nitrogen by heterotrophic bacteria by managing the C:N ratio in the water. Furthermore, the heterotrophic bacteria biomass forms aggregation, which is then called biofloc. Biofloc comprises not only the bacteria itself, but also other microorganism such as microalgae and zooplankton, which are trapped in organic particles or solid. The biofloc technology was good to control both water quality and feed production *in situ*. Biofloc technology offers aquaculture a sustainable tool to simultaneously address its environmental [8]. The biofloc technology was proposed as sustainable solution to catfish culture with high density, which can control water quality under negligible water exchange and sustain healthy culture of catfish. This study was conducted to evaluate the effects of stocking density in intensive aquaculture on blood and production performance of catfish (*Clarias gariepinus*) which were reared with biofloc technology.

2.0 EXPERIMENTAL

2.1 Experimental Design

The intensification experiment was done in one way design. Three level of catfish density i.e. 500 fish per m², 1 000 fish per m², and 1 500 fish per m² were stocked in fibre glass tank (2 × 1 × 1) m³. The catfish were reared in zero-water exchange biofloc-base intensive culture tank for a period of 10 wk. Good water quality was maintained with promotion and development of biofloc through sugar addition during the experiment. Variables measured i.e. fish production, growth, survival rate, feed consumption, feeding efficiency, blood cell counting, bilirubin serum, blood glucose level, glutamate pyruvate aminotransferase (GPT), and glutamate oxaloacetate aminotransferase (GOT). Water quality and fish weight were determined every 2 wk. Blood variables were determined in the end of treatment. Values of variables assayed were presented in mean (± SD). One way ANOVA was used to determine significant difference by SPSS 16.0 program. If there were a significant F-test, subsequent comparisons of treatment means were performed using the Tukey's HSD test. The level significant was set at $P < 0.05$.

2.2 Biofloc

Biofloc were developed in the fibre glass tanks (2 000 L each) during the experiment. Biofloc were produced by inoculating heterotrophic bacteria in the experimental tanks with addition of sugar. The volume of water was initially equilibrated through exchange water between the tanks. Water was added to prevent evaporation. Tanks were aerated and stirred continuously using aquarium pump.


2.3 Fish Stocking and Tanks Management

Eighteen thousand *C. gariepinus* fingerlings (3.18 g ± 0.79 g) were stocked to nine fiberglass tanks (2 × 1 × 1) m³ within biofloc system. Fish were stocked at a density of (500, 1 000 and 1 500) fish per m² in each treatment (three treatments with three replications). The experiment was carried out over a 10 wk period to investigate the effects of stocking density on production and blood performance of catfish (*Clarias gariepinus*) which were reared with biofloc technology.


During experiment, fish were fed to satiation twice per day. Fish were fed with a floating pelleted feed, which contains 30 % of protein, 6 % of fiber, 5 % of fat and 13 % of ash, 13 % of water. The feed were weighed to calculate total food consumption. Water quality in the culture system was monitored every 2 wk for dissolved oxygen (DO), temperature, pH, and turbidity using water quality checker (HORIBA). Total ammonia nitrogen (TAN) was measured using Ammonia Test Kit Microquant 114750, Merck, Jerman. Fish were weighed every 2 wk. Relative growth rate expressed in percentage was calculated by subtracting of the initial weight from the final weight and, dividing this by the initial weight and the total numbers of experimental days. Feed efficiency was calculated by dividing the total final biomass minus the total initial biomass by the total feed fed, and it was expressed as a percentage.

In every 2 wk, five fish were trawled randomly from each stocking density fiberglass tank. Chemical, enzyme and blood cell analysis were performed. Blood sample were collected from the caudal vein using heparinised syringes. Photometric blood cell were calculated using ABX Micros 60. Blood cells counting consist of leukocytes, erythrocytes, hemoglobin, hematocrit and platelet. Whole blood sample was centrifuged (5 min, 5 000 rpm) [1 rpm = 1/60 Hz] prior to hematocrite measurement. The remaining blood was centrifuged at 1 200 rpm for 15 min and the plasma was separated for analysis of glucose, bilirubin and aminotransferase enzyme. Blood glucose concentration was measured by the enzymatic reaction between glucose in blood samples with glucose oxidase and potassium ferric cyanide which produces potassium ferrocyanide. Potassium ferrocyanide was formed with a certain proportion of blood glucose levels. Oxidation of potassium ferrocyanide generates electricity which is


then converted into glucose concentration. Total bilirubin and direct bilirubin were measured by photometric method [9]. The principle of the method is that bilirubin reacted with DSA (diazotized sulfanilic acid) and forms a red color. Absorbance values at λ 546 nm represents direct proportion of the bilirubin concentration. Glucuronid bilirubin which is dissolved in water reacts directly with DSA, whereas indirect bilirubin conjugated with albumin will react with DSA only if there is an accelerator. Thus, $\text{total bilirubin} = \text{direct} + \text{indirect}$ bilirubin. The reaction equation is as follows [9]:



Transaminase enzyme activity consisting of GPT and GOT in the blood serum and catfish sample from each treatment was measured using the kinetic method and UV-photometry method. The principle of measuring the reaction of GPT (ALAT) is as follows:



Whereas GOT (ASAT) measured with principle reaction is as follows:



2.4 Statistical Analysis

In this study, fibreglass tanks were used as experimental unit. All calculations were made using the SPSS system (SPSS 16.0). Data from the effects of stocking density were first analysed using one way analysis of variance (ANOVA) followed by the Tukey's HSD post-hoc test. Homogeneity of variance and normality were tested to meet the assumptions of an ANOVA.

3.0 RESULTS AND DISCUSSION

3.1 Growth and Production Performance of Catfish

The increasing of body weight during the experiment was shown in Figure 1. The average of body weight of catfish decreased with the increasing of stocking density. Conversely, the production of catfish increased with the increasing of fish density. The response of fish production to stocking density of the catfish which were reared with biofloc technology is presented in Table 1. The catfish density of (500, 1 000, and 1 500) fish per m^2 significantly affected fish production ($p < 0.05$) in which the value of catfish

production were (68.09 ± 1.00) kg; (87.41 ± 9.24) kg; and (127.29 ± 16.07) kg, respectively. Previously, study in Senegal sole showed that high stocking density might not alter growth [10]. High stocking density is considered a relevant issue concerning welfare of farmed fish [11] and it is also connected to production parameters when commercial farming conditions are developed, in line with this study stocking density affected to catfish production (Table 1).




Figure 1 Catfish body weight during the experiment

The values of relative growth rate were $(33.24 \pm 2.450) \% \text{ d}^{-1}$; $(21.66 \pm 1.82) \% \text{ d}^{-1}$; and $(21.40 \pm 2.72) \% \text{ d}^{-1}$, each for the stocking density of (500, 1 000, and 1 500) fish per m^2 , respectively. The stocking density significantly affected to the relative growth rate ($P < 0.05$). The increasing of relative growth rate value was due to the total feed consumption. The feed consumption of stocking density treatments of (500, 1 000, and 1 500) fish per m^2 were (63.63 ± 2.70) kg; (96.60 ± 2.49) kg; and (141.00 ± 14.17) kg, respectively. The value of relative growth rate and total feed consumption in stocking density treatment (1 000 and 1 500) fish per m^2 was not significant different (Tukey's HSD, $P > 0.05$), but value of relative growth rate and total feed consumption in stocking density treatment 500 fish per m^2 was significantly difference with that value in stocking density treatment (1 000 and 1 500) fish per m^2 .

On the other hand, the values of feed efficiency decreased with the increasing of stocking density. The values of feed efficiency in stocking density (1 000 and 1 500) fish per m^2 were similar (Tuckey HSD, $P > 0.05$). However, the feed efficiency values of the two treatments were lower than the value of the stocking density 500 fish per m^2 .

The values of survival rate for the stocking density of (500, 1 000, and 1 500) fish per m^2 were $(96.07 \pm 0.21) \%$; $(95.38 \pm 0.50) \%$; and $(94.74 \pm 1.30) \%$, respectively. Survival rate was not affected by stocking densities ($P < 0.05$). The high value of that survival rate caused by proper water quality parameters.

Table 1 Production performance of catfish were rearing with the biofloc technology

Stocking density (fish · m ²)	500 ^a	1 000 ^a	1 500 ^a	P-value
Relative growth rate (% · d ⁻¹)	33.24 ± 2.45 ^a	21.66 ± 1.82 ^b	21.40 ± 2.76 ^b	0.0010
Total feed consumption (kg)	63.63 ± 2.70 ^c	96.60 ± 2.49 ^b	141.00 ± 14.17 ^a	0.0001
Feed efficiencies (%)	107.20 ± 4.73 ^a	90.36 ± 5.53 ^b	90.27 ± 1.65 ^b	0.0120
Biomass production netto (kg)	68.09 ± 1.00 ^b	87.41 ± 9.24 ^b	127.29 ± 16.07 ^a	0.0100
Survival rate (%)	96.07 ± 0.21 ^a	95.38 ± 0.50 ^a	94.74 ± 1.30 ^a	0.2190

Note : ^aValue ± SD (Standard Deviation)

3.2 Water Quality

Water quality criteria were in the suitable range (Table 2). Average and range values of water temperature in stoking density of (500, 1 000, and 1 500) fish per m² were (29.55 °C, range 27.5 °C to 30 °C); (27.72 °C, range 27.3 °C to 28.5 °C); and (27.87 °C, range 27.2 °C to 28.5 °C), respectively. The values of dissolved oxygen concentration were 3.37 mg L⁻¹, range (3.2 to 3.69) mg L⁻¹; (3.36 mg L⁻¹ range (3.22 to 3.47) mg L⁻¹); 3.27 mg L⁻¹ range (2.85 to 3.53) mg L⁻¹, respectively for stocking density of (500, 1 000, and 1 500) fish per m², and pH were (7.54, range 7.35 to 7.8); (7.58, range 7.24 to 7.96); (7.51, range 6.88 to 7.92), respectively for stocking density of (500, 1 000, and 1 500) fish per m². The values of temperature, dissolved oxygen, and pH were within the suitable range for catfish culture. While the values of turbidity were (430.67 mg L⁻¹, range (47.22; 27) mg L⁻¹); (667.79 mg L⁻¹, range (5.09 to 999) mg L⁻¹); and (667.82 mg L⁻¹, range (5.34 to 999) mg L⁻¹), respectively for stocking density of (500, 1 000, and 1 500) fish per m². The values of turbidity increased with the increasing of stocking density. Total ammonia nitrogen (TAN) values were (0.51 mg L⁻¹, range (0.00 to 1.81) mg L⁻¹); (0.51 mg L⁻¹, range (0.00 to 1.81) mg L⁻¹); (1.37 mg L⁻¹, range (0.25 to 3.87) mg L⁻¹), respectively for stocking density of (500, 1 000, and 1 500) fish per m². This value were within safe range for catfish culture. The low level of total ammonia nitrogen was due to immobilization of ammonia to bacteria in biofloc technology. The nitrification processes requires dissolved oxygen for oxidation of total ammonia nitrogen. Oxidating of 1 mg TAN required 4 mg oxygen and 8 mg HCO₃⁻. While dissolved Oxygen values were within safe range for catfish culture. However, continuous vigorous aeration in biofloc technology ensured that oxygen was not limiting.

The biofloc systems lead to loss of buffering capacity and rates of nitrification [5]. The rate of nitrification in biofloc systems was coupled with ammonia immobilization into bacteria. It is reported that nitrification requires approximately 1.4 mg O₂ and 8 mg HCO₃⁻ for oxidizing 1 mg TAN. There are three principal pathways to remove hazardous Nitrogen species in aquaculture: (i) photoautotrophic removal

by algae, (ii) immobilization by heterotrophic bacteria as proteinacious microbial biomass and (iii) chemo-autotrophic oxidation of nitrate by nitrifying bacteria [12]. Although the carbohydrate was added based on TAN:CHO of 1:20 [6], TAN concentrations were elevated, sometimes reaching critical levels in rearing tanks during the experiment. The CHO requirement to minimize TAN needs further evaluation, with addition of sugar.

3.3 Hematological and Blood Biochemical Parameters

The evaluation of hematological and biochemical parameters might be useful for the diagnosis of fish pathologies and physiological status [13]. Alteration of blood cell numbers, as well as biochemical and hormonal status might be indicative of unsuitable environmental conditions (temperature, pH, oxygen concentration) or the presence of stressing factors such as common farm operations. For instance, high stocking density is regarded as a chronic stressor eliciting an increase in plasma cortisol, glucose and osmolality levels [14,15]. The values of blood cell, blood biochemistry, blood glucose, and enzyme serum responses of catfish (*Clarias gariepinus*) to stocking density were presented in Table 3. In the present study, hematocrit, relative levels of circulating leucocytes and glucose levels remained were significantly different in densities treatments (Table 3), suggesting that catfish reared at both 500 and 1 500 fish per m² were not able to adapt to the increasing stocking densities under the particular culture conditions and during the experimental period described in this study. These results are in agreement with that reported in other species, such as gilthead seabream (*Sparus aurata*), rainbow trout and European seabass (*Dicentrarchus labrax*) [16–18]. Furthermore, this hypothesis is supported by the lack of the immunosuppressive state typically observed in chronically stressed fish. When a given stressor is chronic, the immune response shows suppressive effects and therefore the chances of an infection may be enhanced [19].

Table 2 Value of water quality parameters during the experiment

Stocking density (Fish m ²)	42	500	1 000	1 500
Temperature (°C)	Average ± SD	29.55 ± 1.14	27.72 ± 0.42	27.87 ± 0.51
	range	27.5 to 30.7	27.3 to 28.5	27.2 to 28.5
Dissolved Oxygen (mg · L ⁻¹)	Average ± SD	3.37 ± 1.14	3.36 ± 0.42	3.27 ± 0.51
	range	3.2 to 3.69	3.22 to 3.47	2.85 to 3.53
pH	Average ± SD	7.54 ± 0.15	7.58 ± 0.27	7.51 ± 0.39
	range	7.35 to 7.8	7.24 to 7.96	6.88 to 7.92
Turbidity (mg · L ⁻¹)	Average ± SD	430.67 ± 226.70	667.79 ± 513.10	667.82 ± 513.06
	range	47 to 627	5.09 to 999	5.34 to 999
Total Ammonia Nitrogen (mg · L ⁻¹)	Average ± SD	0.51 ± 0.45	0.73 ± 0.64	1.37 ± 1.33
	range	0.00 to 1.03	0.00 to 1.81	0.25 to 3.87

Table 3 Blood performances of catfish (*Clarias gariepinus*) response to different stocking density

Stocking density (fish per m ²)	500 ^a	1 000 ^a	1 500 ^a	P-value
Blood cell parameters				
Total Leucocyte (10 ³ sel per µl)	210.73 ± 0.6 ^a	108.36 ± 0.45 ^b	91.25 ± 0.35 ^c	0.0001
Eritrosit (10 ⁶ sel per µl)	1.83 ± 0.06	1.75 ± 0.20	1.57 ± 0.15	0.1770
Hemoglobine (gr · dl ⁻¹)	6.60 ± 0.26 ^b	7.43 ± 0.08 ^a	6.62 ± 0.12 ^b	0.0200
Hematocrite (%)	23.53 ± 0.50 ^b	26.56 ± 0.20 ^a	23.23 ± 0.23 ^b	0.0001
Trombocyte (10 ³ sel per ul)	1.17 ± 0.29 ^c	16.25 ± 0.28 ^a	13.15 ± 0.14 ^b	0.0001
Blood biochemical parameters				
Glucose (mg · dl ⁻¹)	96.00 ± 2.00 ^a	76.20 ± 0.29 ^b	72.97 ± 0.03 ^c	0.0001
Total Bilirubin (mg · dl ⁻¹)	0.40 ± 0.10 ^a	0.08 ± 0.01 ^b	0.04 ± 0.01 ^b	0.0100
Direct bilirubin (mg · dl ⁻¹)	0.27 ± 0.06 ^a	0.03 ± 0.01 ^b	0.02 ± 0.01 ^b	0.0001
Indirect bilirubin (mg · dl ⁻¹)	0.13 ± 0.06 ^a	0.06 ± 0.01 ^{ab}	0.02 ± 0.02 ^b	0.0210
38 µm enzyme parameter				
SGOT (U · L ⁻¹)	194.33 ± 4.04 ^a	145.04 ± 0.04 ^b	123.05 ± 0.05 ^c	0.0001
SGPT (U · L ⁻¹)	99.00 ± 3.61 ^a	57.02 ± 0.03 ^b	39.02 ± 0.03 ^c	0.0001

Note : ^aValue ± SD

The values of blood cell counting which consists of total leucosit, hemoglobine, hematocrite, and trombocyte were affected by stocking density ($P < 0.05$). On the other hand, the values of eritrocite was not affected by treatments. Surprisingly, several parameters in catfish such as total leucocyte values, blood glucose, bilirubin, SGOT, and SGPT declined in response to increasing stocking density. Fish stocking density has a significant effect on fish physiological responses, and the importance of evaluating density stress [20]. The responses of blood performances indicated that stocking of 1 500 fish per m² lead to healthy fish. The effects of stocking density on welfare African catfish was not negatively influenced by increasing density [1].

4.0 CONCLUSION

Stocking density in intensification catfish culture clearly contributed to the fish production, growth,

feed efficiency and fish health. The biofloc technology can effectively remove total ammonia nitrogen (TAN) and thus prevent accumulation of TAN in the water. Optimum stocking density in catfish culture with biofloc technology was 1 500 fish per m². The stocking density in that system can support maximum catfish production, fish health and proper water quality.

Acknowledgement

The authors thank to Ditlitabmas Ditjen Dikti for the financial support (DIPA-023.04.1.673453/2015).

References

- [1] Van de Nieuwegiesse, P. G., J. Olwo, S. Khong, J. A. J. Vareth, and J. W. Schrama, 2009. Effects of Age and

- [1] Stocking Density on the Welfare of African Catfish, *Clarias fariatus* Burchall. *Aquaculture*. 288: 69-75.
- [2] Hossain, M. A. R., M. C. M. Beveridge, and G. S. Haylor. 1998. The Effects of Density, Light and Shelter on the Growth and Survival of African Catfish (*Clarias fariatus* Burchall, 1822) Fingerlings. *Aquaculture*. 168: 251-258.
- [3] Avnimelech, Y., B. Weber, A. Millstein, B. Hefner, and M. Zorom. 1986. Studies in Circulated Fishponds : Organic Matter Recycling and Nitrogen Transformation. *Aquaculture and Fisheries Management*. 17: 231-242.
- [4] Hargreaves, J. A. 2006. Phytosynthetic Suspended-growth System in Aquaculture. *Aquaculture Engineering*. 34: 344-353.
- [5] Azim, M. E. and D. C. Little. 2008. The Biofloc Technology (BFT) in Indoor Tanks: Water Quality, Biofloc Composition, and Growth and Welfare of Nile Tilapia (*Oreochromis niloticus*). *Aquaculture* 238: 29-35.
- [6] Avnimelech, Y. 1999. Carbon and Nitrogen Ratio as a Control Element in Aquaculture Systems. *Aquaculture*. 176: 227-235.
- [7] Kuhn, D. D., A. L. Lawrence, G. D. Boardman, S. Patnaik, L. Marsh, and G. J. Flick, Jr. 2010. Evaluation of Two Type of Bioflocs Derived from Biological Treatment of Fish Effluent as Feed Ingredients for Pacific White Shrimp, *Litopenaeus setiferus*. *Aquaculture*. 303: 28-33.
- [8] Crab, R., T. Defoirdt, P. Bossier, and W. Verstraete. 2012. Biofloc Technology in Aquaculture: Beneficial Effects and Future Challenges. *Aquaculture*. 356: 351-356.
- [9] Jendrassik, L. and P. Grof. 1938. Colorimetric Method of Determination of Bilirubin. *Biochem Z.* 297: 81-82.
- [10] Andrade, T., A. Afonso, A. Pérez-Jiménez, et al. 2015. Evaluation of Different Stocking Densities in a Senegalese Sole (*Solea senegalensis*) Farm: Implications for Growth, Humoral Immune Parameters and Oxidative Status. *Aquaculture*. 438: 6-11.
- [11] Ashley, P. J., 2007. Fish Welfare: Current Issues in Aquaculture. *Appl. Anim. Behav. Sci.* 104: 199-235.
- [12] Ebeling, J. M., M. B. Timmons, and J. J. Bisogni. 2006. Engineering Analysis of the Stoichiometry of Photoautotrophic, Autotrophic, and Heterotrophic Removal of Ammonia-Nitrogen in Aquaculture Systems. *Aquaculture*. 157: 346-358.
- [13] Maita, M. 2007. Fish Health Assessment. In Nakagawa, H., M. Sato, D. M. Gatlin III (eds.). *Dietary Supplements for the Health and Quality of Cultured Fish*. Cambridge: C. A. B. International.
- [14] Mommsen, T. P., M. M. Vijayan, and T. W. Moon. 1999. Cortisol in Teleosts: Dynamics, Mechanisms of Action, and Metabolic Regulation. *Rev. Fish Biol. Fish.* 9: 211-268.
- [15] Wendelaar Bonga, S. E. 1997. The Stress Response in Fish. *Physiol. Rev.* 7: 591-625.
- [16] Morgan, A. L., K. D. Thompson, N. A. Auchinachie, and H. Migaud. 2008. The Effect of Seasonality on Normal Haematological and Innate Immune Parameters of Rainbow Trout *Oncorhynchus mykiss* L. *Fish Shellfish Immunol.* 25: 791-799.
- [17] Pascoli, F., G. S. Lanzano, E. Negrato, C. Poltronieri, A. Trocino, G. Radaelli, and D. Bertotto. 2011. Seasonal Effects on Hematological and Innate Immune Parameters in Sea Bass *Dicentrarchus labrax*. *Fish Shellfish Immunol.* 31: 1081-1087.
- [18] Tort, L., J. Rotllant, and L. Rovira. 1998. Immunological Suppression in Gilthead Sea Bream *Sparus aurata* of the North West Mediterranean at Low Temperature. *Comp. Biochem. Physiol. A.* 120: 175-179.
- [19] Tort, L. 2011. Stress and Immune Modulation in Fish. *Dev. Comp. Immunol.* 35: 1366-1375.
- [20] Hasenbein, M., N. A. Fangue, J. P. Geist, L. M. Komoroski, and R. E. Connon. 2016. Physiological Stress Biomarkers Reveal Stocking Density Effects in Late Larval Delta Smelt (*Hypomesus transpacificus*). *Aquaculture* 450: 108-115 (In Press).

APPLICATION OF BIOFLOC TECHNOLOGY IN INTENSIVE FARMING AFFECTED PRODUCTION AND BLOOD PERFORMANCES OF THE CATF

ORIGINALITY REPORT

27%

SIMILARITY INDEX

1%

INTERNET SOURCES

27%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

M.E. Azim, D.C. Little. "The biofloc technology (BFT) in indoor tanks: Water quality, biofloc composition, and growth and welfare of Nile tilapia (*Oreochromis niloticus*)", *Aquaculture*, 2008

Publication

6%

2

Wen-chang Liu, Guo-zhi Luo, Li Li, Xiao-yong Wang, Jiao Wang, Nian-nian Ma, Da-chuan Sun, Hong-xin Tan. "Nitrogen Dynamics and Biofloc Composition using Biofloc Technology to Treat Aquaculture Solid Waste Mixed with Distillery Spent Wash", *North American Journal of Aquaculture*, 2016

Publication

2%

3

Gregory D. King, Jacqueline M. Chapman, Jonathan D. Midwood, Steven J. Cooke, Cory D. Suski. "Watershed-Scale Land Use Activities Influence the Physiological Condition of Stream Fish", *Physiological and Biochemical Zoology*,

2%

4

Brett W. Herbert. "Effects of Added Shelter and Stocking Density on Growth of Sleepy Cod *Oxyeleotris lineolatus* in Ponds", Journal of the World Aquaculture Society, 12/2003

Publication

5

Millán-Cubillo, Antonio Francisco, Juan Antonio Martos-Sitcha, Ignacio Ruiz-Jarabo, Salvador Cárdenas, and Juan Miguel Mancera. "Low stocking density negatively affects growth, metabolism and stress pathways in juvenile specimens of meagre (*Argyrosomus regius*, Asso 1801)", Aquaculture, 2016.

Publication

6

Barcellos, L.J.G.. "Hematological changes in jundia (*Rhamdia quelen* Quoy and Gaimard Pimelodidae) after acute and chronic stress caused by usual aquacultural management, with emphasis on immunosuppressive effects", Aquaculture, 20040802

Publication

7

T. L Korkea-aho. "Pseudomonas M162 confers protection against rainbow trout fry syndrome by stimulating immunity", Journal of Applied Microbiology, 04/2012

Publication

1 %

1 %

1 %

1 %

8

Prangnell, David I., Leandro F. Castro, Abdulmehdi S. Ali, Craig L. Browdy, Paul V. Zimba, Susan E. Laramore, and Tzachi M. Samocha. "Some Limiting Factors in Superintensive Production of Juvenile Pacific White Shrimp, *Litopenaeus vannamei* , in No-water-exchange, Biofloc-dominated Systems : SUPERINTENSIVE PRODUCTION OF L. VANNAMEI IN BIOFLOC SYSTEMS", Journal of the World Aquaculture Society, 2016.

Publication

1%

9

Marina Casimiro. "Dietary tryptophan and methionine as modulators of the european seabass (*Dicentrarchus labrax*) immune status and inflammatory response", Repositório Aberto da Universidade do Porto, 2014.

Publication

1%

10

Davis, B. E., N. A. Miller, E. E. Flynn, and A. E. Todgham. "Juvenile Antarctic rockcod (*Trematomus bernacchii*) are physiologically robust to CO₂-acidified seawater", Journal of Experimental Biology, 2016.

Publication

1%

11

Adriana Borges. "Hematologic and serum biochemical values for jundiá (*Rhamdia quelen*)", Fish Physiology and Biochemistry, 03/2004

Publication

1%

12

Mostafa A.R. Hossain, Malcolm C.M. Beveridge, Graham S. Haylor. "The effects of density, light and shelter on the growth and survival of African catfish (*Clarias gariepinus* Burchell, 1822) fingerlings", *Aquaculture*, 1998

Publication

1%

13

E. Gisbert, A. Skalli, J. Campbell, M. M. Solovyev, C. Rodríguez, J. Dias, J. Polo. "Spray-dried plasma promotes growth, modulates the activity of antioxidant defenses, and enhances the immune status of gilthead sea bream () fingerlings", *Journal of Animal Science*, 2015

Publication

1%

14

Haque, M Rezoanul, M Ashraful Islam, M Mojibar Rahman, Mst Farzana Shirin, M Abdul Wahab, and M Ekram Azim. "Effects of C/N ratio and periphyton substrates on pond ecology and production performance in giant freshwater prawn *Macrobrachium rosenbergii* (De Man, 1879) and tilapia *Oreochromis niloticus* (Linnaeus, 1758) polyculture system", *Aquaculture Research*, 2013.

Publication

1%

15

Irkin, Latife Ceyda, Murat Yigit, Sevdan Yilmaz, and Masashi Maita. "Toxicological Evaluation of Dietary Garlic (*Allium sativum*) Powder in

1%

European Sea Bass *Dicentrarchus labrax*
Juveniles", Food and Nutrition Sciences, 2014.

Publication

16

Jeonghwan Park, Luke A. Roy, Nilima Renukdas, Trevor Luna. " Evaluation of a Biofloc System for Intensive Culture of Fathead Minnows, ", Journal of the World Aquaculture Society, 2017

Publication

1 %

17

Toko, I.. "Rearing of African catfish (*Clarias gariepinus*) and vundu catfish (*Heterobranchus longifilis*) in traditional fish ponds (whedos): Effect of stocking density on growth, production and body composition", Aquaculture, 20070214

Publication

<1 %

18

Lima, Ronaldo L. de, Neiva Braun, Daiani Kochhann, Rafael Lazzari, João Radünz Neto, Bibiana S. Moraes, Vania L. Loro, and Bernardo Baldisserotto. "Survival, growth and metabolic parameters of silver catfish, *Rhamdia quelen*, juveniles exposed to different waterborne nitrite levels", Neotropical Ichthyology, 2011.

Publication

<1 %

19

Chhorn Lim. "Growth performance, vitamin E status, and proximate and fatty acid composition of channel catfish, *Ictalurus*

<1 %

punctatus, fed diets containing various levels of fish oil and vitamin E", Fish Physiology and Biochemistry, 10/10/2009

Publication

20

Jith K. Thomas, Oana Birceanu, Bastien Sadoul, Mathilakath M. Vijayan. "Bisphenol A in Eggs Impairs the Long-Term Stress Performance of Rainbow Trout in Two Generations", Environmental Science & Technology, 2018

Publication

<1 %

21

Maurcio Emerenciano, Gabriela Gaxiola, Gerard Cuzo. "Chapter 12 Biofloc Technology (BFT): A Review for Aquaculture Application and Animal Food Industry", InTech, 2013

Publication

<1 %

22

Fayun Li, Ying Zhang, Zaiping Xiong, Tingting Sun. "Effect of Deicing Salt on Ion Concentrations in Urban Roadside Snow and Surface Water", 2011 5th International Conference on Bioinformatics and Biomedical Engineering, 2011

Publication

<1 %

23

Chaudhary, Siddhartha Shankar, Jay Prakash Shah, and Ram Vinod Mahato. "Interference of Bilirubin in Creatinine Value Measurement by Jaffe Kinetic Method", Annals of Clinical

<1 %

24

Jiwyam, Wirat, and Nudtha Nithikulworawong. "Stocking density-dependent growth and survival of Asian red-tailed catfish (*Hemibagrus wyckioides*) fries: early nursing in cages", International Aquatic Research, 2014.

Publication

25

Uloma B. Elvis-Offiah, Enitome E. Bafor, Gerald I. Eze, Osamwonyi Igbinumwen, Christina Viegelmann, RuAngelie Edrada-Ebel. " In vivo investigation of female reproductive functions and parameters in nonpregnant mice models and mass spectrometric analysis of the methanol leaf extract of (Sims) G Dons ", Physiological Reports, 2016

Publication

26

Emmanuelle Roque d'Orbcastel. "Effects of rearing density on sea bass (*Dicentrarchus labrax*) biological performance, blood parameters and disease resistance in a flow through system", Aquatic Living Resources, 2009

Publication

27

Fagbenro. "Use of soybean flour (dehulled, solvent-extracted soybean) as a fish meal substitute in practical diets for African catfish,

<1 %

<1 %

<1 %

<1 %

Clarias gariepinus (Burchell 1822): growth, feed utilization and digestibility", Journal of Applied Ichthyology, 4/28/2001

Publication

28

Catarina I. M. Martins. "Behavioural indicators of welfare in farmed fish", Fish Physiology and Biochemistry, 07/28/2011

Publication

<1 %

29

Chiang, P.D.. "Non-volatile taste components of various broth cubes", Food Chemistry, 2007

Publication

<1 %

30

Mohamd Moaaz Refa, Ahmed Ismail Meh, Magda Mahmoud El. "Aquafat-O® as a Growth and Physiological Enhancer Agent for Hybrid Red Tilapia, ♂Oreochromis niloticus (L.)×♀Oreochromis mossambicus (Peters)", Asian Journal of Animal and Veterinary Advances, 2017

Publication

<1 %

31

Kuhn, David D., Addison L. Lawrence, Jack Crockett, and Dan Taylor. "Evaluation of bioflocs derived from confectionary food effluent water as a replacement feed ingredient for fishmeal or soy meal for shrimp", Aquaculture, 2016.

Publication

<1 %

32

Eglal Ali Omar. "Optimum dietary protein level

and stocking density for catfish (*Clarias lazera*) fingerlings in concrete ponds", Journal of Animal Physiology and Animal Nutrition, 09/12/1996

Publication

<1 %

33

Crab, Roselien, Tom Defoirdt, Peter Bossier, and Willy Verstraete. "Biofloc technology in aquaculture: Beneficial effects and future challenges", Aquaculture, 2012.

Publication

<1 %

34

Caipang, Christopher Marlowe A., Han Xuan Choo, Zemin Bai, Huilin Huang, and Clara M. Lay-yag. "Viability of sweet potato flour as carbon source for the production of biofloc in freshwater culture of tilapia, *Oreochromis* sp.", International Aquatic Research, 2015.

Publication

<1 %

35

fishethobase.fair-fish.net

Internet Source

<1 %

36

Ziad, . "Impact of Shrimp Farm Effluent on Water Quality in Coastal Areas of the World Heritage-Listed Ha Long Bay", American Journal of Environmental Sciences, 2012.

Publication

<1 %

37

Harsha Haridas, Ajit Kumar Verma, Gaurav Rathore, Chandra Prakash, Paramita Banerjee Sawant, Asanaru Majeedkutty Babitha Rani.

<1 %

"Enhanced growth and immuno-physiological response of Genetically Improved Farmed Tilapia in indoor biofloc units at different stocking densities", Aquaculture Research, 2017

Publication

38

Narayanaperumal Pravin, Ganesan Kumaravel, Raju Senthilkumar, Natarajan Raman. " Water-soluble Schiff base Cu(II) and Zn(II) complexes: Synthesis, DNA targeting ability and chemotherapeutic potential of Cu(II) complex for hepatocellular carcinoma - and approach ", Applied Organometallic Chemistry, 2017

<1 %

Publication

39

"Wastewater Treatment, Plant Dynamics and Management in Constructed and Natural Wetlands", Springer Nature America, Inc, 2008

<1 %

Publication

40

G. S. HAYLOR. "Controlled hatchery production of *Clarias gariepinus* (Burchell 1822): growth and survival of fry at high stocking density", Aquaculture Research, 12/1991

<1 %

Publication

41

Amal Biswas, Kenji Takii. " Optimal Stocking Density with the Provision of Self-feeders for Striped Knifejaw, (Temminck and Schlegel 1844) ", Journal of the World Aquaculture

<1 %

-
- 42 Zhan-Jun Xue. "Health risk assessment of heavy metals for edible parts of vegetables grown in sewage-irrigated soils in suburbs of Baoding City, China", Environmental Monitoring and Assessment, 07/16/2011

Publication

-
- 43 Özgür Fırat. "Individual and Combined Effects of Heavy Metals on Serum Biochemistry of Nile Tilapia *Oreochromis niloticus*", Archives of Environmental Contamination and Toxicology, 06/02/2009

Publication

-
- 44 G.M. Naja, B. Volesky. "Constructed Wetlands for Water Treatment", Elsevier BV, 2011

Publication

-
- 45 Wang, Guangjun, Ermeng Yu, Jun Xie, Deguang Yu, Zhifei Li, Wen Luo, Lijiang Qiu, and Zonglin Zheng. "Effect of C/N ratio on water quality in zero-water exchange tanks and the biofloc supplementation in feed on the growth performance of crucian carp, *Carassius auratus*", Aquaculture, 2015.

Publication

-
- 46 Kuhn, D.D.. "Evaluation of two types of bioflocs derived from biological treatment of fish

effluent as feed ingredients for Pacific white shrimp, *Litopenaeus vannamei*", Aquaculture, 20100524

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

APPLICATION OF BIOFLOC TECHNOLOGY IN INTENSIVE FARMING AFFECTED PRODUCTION AND BLOOD PERFORMANCES OF THE CATF

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6